


Rosenbluth Separation of electropion production cross-section from Hydrogen and Carbon

Xin Qian

Duke University, Durham, NC, USA


On the behalf of Jefferson Laboratory E01-107 collaboration

- Introduction
- Overview of E01-107
- Preliminary results
- Summary



Motivation of E01-107

- ❖ Search for Color-Transparency
- ❖ **Colour Transparency** is a phenomenon predicted by QCD in which hadrons produced at large momentum transfer can pass through nuclear matter with little or no interaction
- $\bar{q}q$ or qqq that have small transverse size are preferentially selected at large Q^2 (**Quantum mechanics**)
- The hadron can propagate out of the nucleus before returning to its equilibrium size (**Relativity**)
- Reduced interaction, $\sigma_{\text{PLC}} \propto (r_{\text{PLC}})^2$



Motivation of Rosenbluth Separation

❖ Measuring Nuclear Transparency

∞ Nuclear Transparency is defined by :

$$R_T = \frac{Y_{data}^{nucleus} / Y_{SIMC}^{nucleus}}{Y_{data}^{hydrogen} / Y_{SIMC}^{hydrogen}}$$

❖ Expected Yield can be calculated used realistic nucleon momentum distributions under quasi-free assumption.

∞ Quasi-free assumption can be verified by carrying out Rosenbluth separation.

$$\frac{\sigma_L^{hydrogen}}{\sigma_T^{hydrogen}} \sim \frac{\sigma_L^{nucleus}}{\sigma_T^{nucleus}}$$

Kinematics

- ❖ Electro pion five-fold DXs can be written as:

$$\frac{d^5\sigma}{d\Omega_e dE_e d\Omega_\pi} = \Gamma \frac{d^2\sigma}{d\Omega_\pi}$$

Γ : virtual photon flux.

- ❖ Photo pion DXs can be decomposed by virtual photon polarization:

$$\frac{d^2\sigma}{d\Omega_\pi} = \epsilon \frac{d^2\sigma_L}{d\Omega_\pi} + \frac{d^2\sigma_T}{d\Omega_\pi} + \sqrt{2\epsilon(\epsilon+1)} \frac{d^2\sigma_{LT}}{d\Omega_\pi} \cos(\phi_\pi) + \epsilon \frac{d^2\sigma_{TT}}{d\Omega_\pi} \cos(2\phi_\pi)$$

ϵ : virtual photon polarization






FIGURE 1. The $(e, e'\pi)$ reaction in lab frame.

In parallel kinematics ($\theta_\pi = 0$)



Rosenbluth Separation

- ❖ By performing experiment at two values of virtual photon polarization, we can extract longitudinal and transverse electro-pion production DXs:

$$\left\langle \frac{d^2\sigma_L}{d\Omega_\pi} \right\rangle = \frac{\left\langle \frac{d^2\sigma_1}{d\Omega_\pi} \right\rangle - \left\langle \frac{d^2\sigma_2}{d\Omega_\pi} \right\rangle}{\left\langle \epsilon_1 \right\rangle - \left\langle \epsilon_2 \right\rangle}$$

$$\left\langle \frac{d^2\sigma_T}{d\Omega_\pi} \right\rangle = \frac{\left\langle \frac{d^2\sigma_1}{d\Omega_\pi} \right\rangle \cdot \left\langle \epsilon_2 \right\rangle - \left\langle \frac{d^2\sigma_2}{d\Omega_\pi} \right\rangle \cdot \left\langle \epsilon_1 \right\rangle}{\left\langle \epsilon_1 \right\rangle - \left\langle \epsilon_2 \right\rangle}$$

Overview of E01-107


- Spokespersons: *D. Dutta, R. Ent* and *K. Garrow*
- Experiment ran at Jefferson Lab in Hall C in 2004
- Standard Hall C equipment was used

Electron beam energy
(4.0 to 5.8 GeV)

Electron in the SOS
(0.73 to 1.73 GeV/c)

Pion in the HMS
(2.1 to 4.4 GeV/c)

$(e, e' \pi^+)$



Kinematics

LH_2 , LD_2 , ^{12}C , ^{63}Cu and ^{197}Au targets at each kinematic setting

	Q^2 (GeV ²)	W (GeV)	-t (GeV ²)	E _{beam} (GeV)	θ _{hms} (deg)	P _{hms} (GeV/c)	θ _{sos} (deg)	P _{sos} (GeV/c)	x _{BJ}	
L-T separation L-T separation W vs k _π test point	1.1	2.3	0.05	4.0	10.6	2.8	27.8	-1.2	0.50	0.21
	2.15	2.2	0.16	5.0	13.4	3.2	28.9	-1.7	0.56	0.35
	3.0	2.1	0.29	5.0	12.7	3.4	37.8	-1.4	0.45	0.44
	4.0	2.2	0.40	5.8	11.5	4.1	40.4	-1.5	0.39	0.50
	4.8	2.2	0.52	5.8	10.6	4.4	52.7	-1.1	0.26	0.54
	2.15	2.2	0.16	4.0	10.6	3.2	50.8	-0.7	0.27	0.35
	4.0	2.1	0.44	5.0	10.6	3.9	55.9	-0.9	0.25	0.52
	2.15	1.7	0.37	4.0	20.0	2.1	32.3	-1.7	0.63	0.50


(k_{π} = momentum of the virtual pion)

Particle Identification (PID)


Electron arm
(SOS) at 1.4 GeV

Cerenkov effic =
99.4%

No Cerenkov cut




One P.E. Cerenkov cut




Pion arm (HMS)
at 3.2 GeV

Cerenkov effic =
98.5%


No Cerenkov cut



0.7 P.E. Cerenkov cut




Preliminary results at $Q^2 = 2.15 \text{ GeV}^2$



statistical uncertainties only


Carbon points have been shifted by 0.01 for clarity

Preliminary results at $Q^2 = 2.15 \text{ GeV}^2$



statistical uncertainties only
Carbon points have been shifted by 0.01 for clarity


Preliminary results at $Q^2 = 4.0 \text{ GeV}^2$



statistical uncertainties only

Carbon points have been shifted by 0.01 for clarity


Preliminary results at $Q^2 = 4.0 \text{ GeV}^2$



statistical uncertainties only

Carbon points have been shifted by 0.01 for clarity

Preliminary results on ratio



statistical uncertainties only

Carbon points have been shifted by 0.1 for clarity

Summary

- **E01-107 will provide the FIRST nuclear transparency data from $(e,e \pi^+)$ reactions.**
- **Rosenbluth separation has been carried out for the first time with $(e,e' \pi^+)$ on Carbon at $Q^2 = 2.15$ and 4.0 GeV^2 and Hydrogen at $Q^2 = 4.0 \text{ GeV}^2$.**
- **Preliminary results are in good agreement with quasi-free assumptions for $Q^2 = 2.15$ and 4.0 GeV^2 .**
- **Rosenbluth separation for Copper and Gold targets will be carried out in the near future.**

E01-107 collaboration

Y. Liang

American University, Washington, DC

J. Arrington, L. El Fassi, X. Zheng

Argonne National Laboratory, Argonne, IL

T. Mertens, D. Rohe

Basel University, Basel, Switzerland

R. Monson

Central Michigan University, Mount Pleasant, MI

C. Perdrisat

College of William and Mary, Williamsburg, VA

D. Dutta (Spokesperson), H. Gao, K. Kramer, X. Qian

Duke University, Durham, NC

W. Boeglin, P. Markowitz

Florida International University, Miami, FL

M. E. Christy, C. E. Keppel, S. Malace, E. Segbefia, L. Tang, L. Yuan

Hampton University, Hampton, VA

J. Ferrer, G. Niculescu, I. Niculescu

James Madison University, Harrisonburg, VA

P. Bosted, A. Bruell, R. Carlini, E. Chudakov, V. Dharmawardane,

R. Ent (Spokesperson), H. Fenker, D. Gaskell, M. K. Jones, A. Lung,

D. G. Meekins, G. Smith, W. F. Vulcan, S. A. Wood

Jefferson Laboratory, Newport News, VA

B. Clisie, J. Seely

Massachusetts Institute of Technology, Cambridge, MA

V. Punjabi

Norfolk State University, Norfolk, VA

A. K. Opper

Ohio University, Athens, OH

A. Villano

Rensselaer Polytechnic Institute, Troy, NY

F. Benmokhtar

Rutgers University, Piscataway, NJ and

Universite' des Sciences et de la Technologie, Algiers, Algeria

Y. Okayasu, A. Matsumura, T. Miyoshi, M. Sumihama

Tohoku University, Sendai, Japan

K. Garrow (Spokesperson)

TRIUMF, Vancouver, British Columbia, Canada

A. Daniel, N. Kalantarians, Y. Li, V. Rodriguez

University of Houston, Houston, TX

A. W. Rauf

University of Manitoba, Winnipeg, Manitoba, Canada

T. Horn

University of Maryland, College Park, MD

G. M. Huber

University of Regina, Regina, Saskatchewan, Canada

D. Day, N. Fomin

University of Virginia, Charlottesville, VA

M. Dalton, C. Gray

University of the Witwatersrand, Johannesburg, South Africa

R. Asaturyan, H. Mkrtchyan, T. Navasardyan, V. Tadevosyan

Yervan Physics Institute, Yervan, Armenia

Systematic uncertainty estimation

❖ SOS Cerenkov Efficiency	0.5 (pt to pt)	❖ Kinematics spcentral	0.5
❖ HMS Cerenkov Efficiency	1.0 (pt to pt)	❖ Kinematics hstheta	0.3
❖ Charge	1.0 (Normalization)	❖ Kinematics hpcentral	0.3
❖ Target thickness	1.0 (Normalization)	❖ Pion decay	2.0 (Pt to Pt)
❖ HMS and SOS trigger efficiency	2.0 (Pt to Pt)	❖ Collimator punch-through	3.0 (Pt to Pt)
❖ Computer dead time	0.1	❖ Radiative correction	2.5 (Pt to Pt)
❖ Coincidence blocking	0.1	❖ Acceptance	5.0 (Pt to Pt)
❖ Tracking efficiency	0.5	❖ Dummy subtraction	0.2 (Pt to Pt)
❖ Pion absorption	3.0	❖ HMS electronic dead time	0.4
(normalization)		❖ SOS electronic dead time	0.3
❖ Pion absorption (between target)	1.0	❖ Target boiling	1.0 (Normalization)
❖ Kinematics Ebeam	0.5	❖ Carbon spectral function	1.0 (Normalization)
❖ Kinematics sstheta	0.5	❖ Model dependence	10.0

Hydrogen DXs: 7.99 %

Carbon DXs: 12.84 %

The estimated systematic uncertainties at this stage are 7% pt-pt, 3.6% normalization and 10% model dependent. We expect to improve several of these uncertainties.